

Complete sets need not be reduced in Minkowski spaces

Horst Martini · Senlin Wu

Received: date / Accepted: date

Abstract It is well known that in n -dimensional Euclidean space ($n \geq 2$) the classes of (diametrically) complete sets and of bodies of constant width coincide. Due to this, they both form a proper subfamily of the class of reduced bodies. For n -dimensional Minkowski spaces, this coincidence is no longer true if $n \geq 3$. Thus, the question occurs whether for $n \geq 3$ any complete set is reduced. Answering this in the negative for $n \geq 3$, we construct $(2^k - 1)$ -dimensional ($k \geq 2$) complete sets which are not reduced.

Keywords bodies of constant width · (diametrically) complete sets · Hadamard matrix · Minkowski Geometry · reduced body · Walsh matrix

Mathematics Subject Classification (2000) 46B20 · 52A20 · 52A21 · 52A40 · 52B11

1 Introduction

We denote by $X = (\mathbb{R}^n, \|\cdot\|)$ an n -dimensional normed or *Minkowski space*, i.e., an n -dimensional real Banach space with origin o . We will use the common abbreviations *aff* and *conv* for affine and convex hull, respectively. A *convex body* in X is a compact, convex set with interior points, and the norm $\|\cdot\|$ is induced by the o -symmetric convex body B_X , called the *unit ball* of X , via

$$\|x\| = \min\{\lambda \geq 0 : x \in \lambda B_X\}, \quad \forall x \in \mathbb{R}^n.$$

Senlin Wu is partially supported by the National Natural Science Foundation of China (grant numbers 11371114 and 11171082) and by Deutscher Akademischer Austauschdienst.

Horst Martini
Faculty of Mathematics, Chemnitz University of Technology, 09107 Chemnitz, Germany
E-mail: horst.martini@mathematik.tu-chemnitz.de

Senlin Wu
Department of Applied Mathematics, Harbin University of Science and Technology, 150080 Harbin, China
E-mail: wusenlin@outlook.com

An interesting part of the geometry of Minkowski spaces (see [9], [4], and [3]) refers to natural extensions of notions from classical convexity to normed spaces. Such notions are, e.g., special classes of convex bodies, like those of constant width, complete sets, and reduced bodies. It is well known that also in Minkowski spaces the *width* of a convex body K in direction u is the (Minkowskian) distance of the two supporting hyperplanes H_1, H_2 of K which are both parallel to the $(n - 1)$ -dimensional subspace orthogonal (in the Euclidean sense) to u . The minimum and the maximum of the *width function* of K are called the *thickness* $\Delta(K)$ and the *diameter* $\delta(K)$ of K , respectively. In any Minkowski space X , *bodies of constant width* are precisely determined by the condition $\Delta(K) = \delta(K)$, and a convex body is called (*diametrically*) *complete* in X if it is not properly contained in a compact set of the same diameter. In a sense dually, a convex body R is said to be *reduced* in X if $\Delta(C) < \Delta(R)$ holds for any convex body C properly contained in R . There are interesting open problems about complete and reduced bodies in Minkowski spaces (see the survey [2]), and even for the Euclidean norm surprisingly elementary questions are still unsolved (cf. [1]). E.g., it is not known whether in Euclidean n -space E^n ($n \geq 3$) there are reduced convex n -polytopes.

It is also interesting that the notion of reduced body, although it somehow dualizes that of completeness, yields a proper superset of the class of complete sets in E^n . The reason is that in E^n the notions of constant width and completeness coincide, and that one can easily construct reduced bodies that are not of constant width. In Minkowski spaces, this coincidence of constant width and completeness is only assured for $n = 2$ (Meissner's theorem; see p. 98 in [3]), and therefore in any Minkowski plane the class of reduced bodies clearly contains that of complete sets. Thus, the question remains whether in n -dimensional Minkowski spaces ($n \geq 3$) any complete set is reduced. A wrong short formulation was given in the survey [2] (cf. line 10 of the Abstract and, in repeated form, line 13 of page 406), which would imply that in Minkowski spaces any complete set is reduced. To correct this, we want to construct non-planar complete sets that are not reduced.

2 Results

Let X be the Minkowski space l_1^3 , i.e., the Banach space on \mathbb{R}^3 endowed with the taxicab norm. Then B_X is the crosspolytope whose vertices are

$$\pm(1, 0, 0), \pm(0, 1, 0), \text{ and } \pm(0, 0, 1).$$

Put K to be the convex hull of the following four points (see Figure 1):

$$a_1 = (-1, -1, -1), a_2 = (1, 1, -1), a_3 = (1, -1, 1), \text{ and } a_4 = (-1, 1, 1).$$

Clearly,

$$\|a_i - a_j\| = 4, \forall \{i, j\} \subseteq \{1, 2, 3, 4\}. \quad (1)$$

Claim 1: The tetrahedron K is a complete set whose diameter is 4.

Proof We only need to show that, for each boundary point x of K , there exists a point $y \in K$ such that $\|x - y\| = \delta(K)$, which follows directly from (1) and the trivial fact that

$$\left\| a_i - \frac{1}{3}(a_j + a_k + a_l) \right\| = 4, \forall \{i, j, k, l\} = \{1, 2, 3, 4\}.$$




Fig. 1 The tetrahedron K is not reduced.

□

Claim 2: $\Delta(K) = 2$.

Proof From the equalities

$$\begin{aligned} \frac{1}{2}(a_2 + a_3) &= (1, 0, 0), & \frac{1}{2}(a_1 + a_2) &= (0, 0, -1), \\ \frac{1}{2}(a_2 + a_4) &= (0, 1, 0), & \frac{1}{2}(a_1 + a_3) &= (0, -1, 0), \\ \frac{1}{2}(a_3 + a_4) &= (0, 0, 1), & \frac{1}{2}(a_1 + a_4) &= (-1, 0, 0), \end{aligned}$$

it follows that $B_X \subseteq K$. Therefore $\Delta(K) \geq 2$. Since the hyperplanes

$$\{(\alpha, \beta, \gamma) : \gamma = 1\} \text{ and } \{(\alpha, \beta, \gamma) : \gamma = -1\}$$

are two parallel supporting hyperplanes of K and the distance between them is 2, $\Delta(K) \leq 2$. Therefore, $\Delta(K) = 2$. □

Now we are ready to show that K is not reduced. Namely, the thickness of the convex body

$$K \cap \{(\alpha, \beta, \gamma) : \alpha + \beta + \gamma \geq -1\},$$

which is a proper subset of K , is still 2.

Now we try to extend this example to higher dimensions. Our main tool is that of Walsh matrices $\mathbf{H}(2^n)$ (cf. [10, Chapters I and II] and [8]) which can be constructed by the following procedure:

$$\begin{aligned} \mathbf{H}(2^1) &= \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \\ \mathbf{H}(2^2) &= \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}, \\ \mathbf{H}(2^k) &= \begin{pmatrix} H(2^{k-1}) & H(2^{k-1}) \\ H(2^{k-1}) & -H(2^{k-1}) \end{pmatrix}, \quad \forall k \in \mathbb{N}, k \geq 2. \end{aligned}$$

Walsh matrices and the related Hadamard matrices can be a helpful tool for modelling and solving geometric problems in Banach spaces and high dimensions (see, e.g., [7] and [11, p. 51]). Since any Walsh matrix is a Hadamard matrix, we have

$$\mathbf{H}(2^n)\mathbf{H}(2^n)^T = 2^n \mathbf{I}_{2^n}, \quad \forall n \in \mathbb{N}, n \geq 1. \quad (2)$$

For each integer $n \geq 2$, let $(\alpha_{i,j})_{0 \leq i,j \leq 2^n-1}$ be the matrix $\mathbf{H}(2^n)$. For each integer $0 \leq i \leq 2^n - 1$, put

$$b_i = (\alpha_{i,j})_{j=0}^{2^n-1} \text{ and } a_i = (\alpha_{i,j})_{j=1}^{2^n-1}.$$

Then

$$V = \{a_i : 0 \leq i \leq 2^n - 1\}$$

is a set of 2^n points in \mathbb{R}^{2^n-1} . By (2), we have that

$$\{b_i : 0 \leq i \leq 2^n - 1\}$$

is linearly independent and, therefore, V is affinely independent. Thus $S = \text{conv} V$ is a simplex in \mathbb{R}^{2^n-1} .

Proposition 3 *Let $X = l_1^{2^n-1}$, and $\|\cdot\|$ be the norm of X . Then*

1. $\delta(S) = 2^n$;
2. for each vertex v of S and each point w in the facet of S opposite to v , $\|v - w\| = 2^n$; moreover, the distance from v to $\text{aff} F_v$ is 2^n ;
3. $B_X \subseteq S$;
4. $\Delta(S) = 2$;
5. S is not reduced.

Proof 1. By (2), for each set $\{i, j\} \subset \{k : k \in \mathbb{Z}, 0 \leq k \leq 2^n - 1\}$ the inner product of b_i and b_j is 0. Therefore b_i and b_j are different in precisely 2^{n-1} coordinates. This implies that

$$\|a_i - a_j\| = 2^n.$$

Thus, the distance between each pair of the vertices of S is 2^n , which implies that $\delta(S) = 2^n$.

2. Being a Hadamard matrix, $\mathbf{H}(2^n)$ must have the following property: each column of $\mathbf{H}(2^n)$, except for the first one, is evenly divided between 1 and -1 . Thus

$$\sum_{i=0}^{2^n-1} a_i = o.$$

Let v be an arbitrary vertex of S , and F_v be the facet of S opposite to v . Then $F_v = \text{conv}(V \setminus \{v\})$, and the point

$$w = \frac{1}{2^n - 1} \sum_{z \in V \setminus \{v\}} z$$

is a relatively interior point of F_v . Moreover,

$$\begin{aligned}\|v - w\| &= \left\| v - \frac{1}{2^n - 1} \sum_{z \in V \setminus \{v\}} z \right\| \\ &= \left\| v + \frac{1}{2^n - 1} v \right\| \\ &= 2^n.\end{aligned}$$

Since the functional

$$\begin{aligned}f : F_v &\mapsto \mathbb{R} \\ z &\mapsto \|v - z\|\end{aligned}$$

is convex,

$$\|v - z\| = 2^n, \quad \forall z \in F_v.$$

The second part of the result follows from the fact that the functional

$$\begin{aligned}f : \text{aff} F_v &\mapsto \mathbb{R} \\ z &\mapsto \|v - z\|\end{aligned}$$

is convex and constant in F_v .

3. For each vertex v of S , denote by H_v the supporting halfspace of S determined by F_v . Then

$$S = \bigcap_{v \in V} H_v.$$

Thus we only need to show that, for each $v \in V$, H_v is also a supporting halfspace of B_X . Since $o \in S$, as can be easily verified, we only need to show that each hyperplane determined by F_v , which is $\text{aff} F_v$, is a supporting hyperplane of B_X at the point $-(1/(2^n - 1))v$. Clearly,

$$-\frac{1}{2^n - 1}v \in F_v \cap B_X.$$

We only deal with the case when $v = a_0$; the other cases can be dealt with in a similar way. For each point $w \in \text{aff} F_v$, there exist numbers $\lambda_1, \dots, \lambda_{2^n - 1}$ such that

$$w = \sum_{i=1}^{2^n - 1} \lambda_i a_i \quad \text{and} \quad \sum_{i=1}^{2^n - 1} \lambda_i = 1.$$

Therefore,

$$\begin{aligned}\|w\| &= \frac{1}{2^n} \|2^n w + v - v\| \\ &= \frac{1}{2^n} \left\| w + (2^n - 1) \left(w + \frac{1}{2^n - 1} v \right) - v \right\| \\ &\geq \frac{1}{2^n} 2^n = 1,\end{aligned}$$

and the inequality holds since the distance from v to $\text{aff} F_v$ is 2^n and

$$w + (2^n - 1) \left(w + \frac{1}{2^n - 1} v \right) = w + (2^n - 1) \left(w - \frac{-1}{2^n - 1} v \right) \in \text{aff} F_v.$$

Thus $\text{aff} F_v$ is a supporting hyperplane of B_X . It follows that $B_X \subseteq S$.

4. Put

$$H_1 = \{(\eta_i)_{i=1}^{2^n-1} : \eta_{2^n-1} = 1\}, \quad H_2 = \{(\eta_i)_{i=1}^{2^n-1} : \eta_{2^n-1} = -1\}.$$

We have $B_X \subseteq S$. Thus, to show that $\Delta(S) = 2$, it suffices to show that H_1 and H_2 are supporting hyperplanes of S , which is clear since each of H_1 and H_2 supports both B_X and its polar body

$$Q = \{(\eta_i)_{i=1}^{2^n-1} : \max\{|\eta_i| : 1 \leq i \leq 2^n - 1\} = 1\},$$

and $B_X \subseteq S \subseteq Q$.

5. Let

$$H = \left\{ (\eta_i)_{i=1}^{2^n-1} : \sum_{i=1}^{2^n-1} \eta_i \leq 1 \right\}.$$

Then $S \cap H$, a proper subset of S , is a convex body containing B_X whose thickness is 2. Thus S is not reduced. \square

Proposition 3 also shows that, for any positive number ε , there exists a positive integer n , an n -dimensional Minkowski space X , and a complete set K in X such that the ratio of $\Delta(K)$ to $\delta(K)$ is less than ε . Anyway, since each non-trivial complete set (i.e., a complete set containing at least two distinct points) in a Minkowski space has an interior point, its thickness is strictly greater than 0.

The situation in infinite dimensional Banach spaces is quite different: it is possible that a non-trivial complete set is contained in a hyperplane (cf. [5, Example 4.6]), and the thickness of such a complete set is clearly 0.

We finish this paper with a very natural question, clearly not posed in [2] and referring to all dimensions $n \geq 3$ (by Meissner's theorem the planar case is clear; see [3, p. 98]).

Problem: *In which Minkowski spaces is any complete set also reduced? Try to give geometric characterizations of their unit balls!*

In some sense vice versa, it might be interesting to collect (e.g., as partial results in that direction) typical geometric properties shared by those Minkowski spaces which contain complete sets that are not reduced. All this is directly motivated by our construction.

More generally, it would be nice to give a complete clarification of the containment and intersection relations between the families of reduced, complete, and constant width bodies (and between suitable subfamilies of them) in Minkowski Geometry, for example via a completely described Venn diagram. This is of course closely related to recent research on those Minkowski spaces in which completeness and constant width are equivalent notions; see, e.g., [6].

References

1. Lassak, M., Martini, H.: Reduced convex bodies in Euclidean space - a survey. *Expositiones Math.* **29**, 204–219 (2011)
2. Lassak, M., Martini, H.: Reduced convex bodies in finite dimensional normed spaces: a survey. *Results Math.* **66**, 405–426 (2014)
3. Martini, H., Swanepoel, K.J.: The geometry of Minkowski spaces - a survey, part II. *Expositiones Math.* **22**, 93–144 (2004)
4. Martini, H., Swanepoel, K.J., Weiss, G.: The geometry of Minkowski spaces - a survey, part I. *Expositiones Math.* **19**, 97–142 (2001)
5. Moreno, J.P., Papini, P.L., Phelps, R.R.: Diametrically maximal and constant width sets in Banach spaces. *Canad. J. Math.* **58**(4), 820–842 (2006)
6. Moreno, J.P., Schneider, R.: Structure of the space of diametrically complete sets in a Minkowski space. *Discrete Comput. Geom.* **48**(2), 467–486 (2012)
7. Schütt, C.: Unconditionality in tensor products. *Israel J. Math.* **31**(3–4), 209–216 (1978)
8. Seberry, J., Yamada, M.: Contemporary Design Theory: A Collection of Surveys, chap. Hadamard matrices, sequences, and block designs, pp. 431–560. Wiley Intersci. Ser. Discrete Math. Optim. Wiley, New York (1992)
9. Thompson, A.C.: Minkowski Geometry. Cambridge University Press, Cambridge (1996)
10. Yang, Y.: Theory and Applications of Higher-Dimensional Hadamard Matrices. Kluwer Academic Publishers, Dordrecht and Beijing (2001)
11. Zong, C.: The Cube: A Window to Convex and Discrete Geometry. Cambridge University Press, Cambridge (2006)